Grass and Forage Science (1996) Volume 51, 470–472

Author Index

ADRIAENS F. *see* PHIPPS R.H.
AKKHARATH I., GUPTA M.L. and TULLBERG J.N. The influence of weather on the effectiveness of mechanical and chemical conditioning on drying rate of lucerne hay 96

BAKER L.J. *see* FISHER G.E.J.
BAKEWELL E.L. *see* DAVIES D.R.
BARDGETT R. *see* DAVIES A.
BEATTIE M.M. *see* HEPP C.
BESGA G. *see* ESTAVILLO J.M.
BINNIE R.C., JOHNSTON D.T. and CHESTNUTT D.M.B. The effect of a high-magnesium perennial ryegrass variety on the magnesium status of sheep 456
BRIGNALL P. *see* DAVIES A.
BUCKINGHAM H. *see* SMITH R.S.
BULLARD M.J. *see* SMITH R.S.

CHAPPLE D.G. *see* HAIGH P.M.
CHESTNUTT D.M.B. *see* BINNIE R.C.
CLAWSON S. *see* MOWAT D.J.
COLGROVE P.M. *see* DEL POZO M.
COLLINS R.P., FOTHERGILL M. and RHODES I. Interactions between seedlings of perennial ryegrass and white clover cultivars in establishing swards 163
COLLINS R.P., FOTHERGILL M., POTTER J.F. and RHODES I. The use of repeated-measurements analysis to describe the overwintering of white clover cultivars 116
CORKHILL P. *see* SMITH R.S.
CUSHNAHAN A. *see* MCKEE C.A.

DAVIES A., SHAO J., BRIGNALL P., BARDGETT R., PARRY M. and POLLOCK C.J. Specification of climatic sensitivity of forage maize to climate change 306
DAVIES D.R., MERRY R.J. and BAKEWELL E.L. The effect of timing of slurry application on the microflora of grass, and changes occurring during silage fermentation 42
DE KERCHOVE G. *see* PHIPPS R.H.
DEL POZO M., WRIGHT I.A., WHYTE T.K. and COLGROVE P.M. Effects of grazing by sheep or goats on sward composition in ryegrass/white clover pasture and on subsequent performance of weaned lambs 142
DOWDESWELL A.M. *see* FISHER G.E.J.

EDWARDS G.R., PARSONS A.J., NEWMAN J.A. and WRIGHT I.A. The spatial pattern of vegetation in cut and grazed grass/white clover pastures 227, 219
ELBERSON H.W. and WEST C.P. Growth and water relations of field-grown tall fescue as influenced by drought and endophyte 333
ESTAVILLO J.M., GONZALEZ-MURUA C., BESGA G. and RODRIGUEZ M. Effect of cow slurry N on herbage productivity, efficiency of N utilization and on white clover content in a natural sward in the Basque Country, Spain 1
ESTELL R.E., FREDRICKSON E.L. and HAVSTAD K.M. Chemical composition of *Flourensia cernua* at four growth stages 434

EVANS D.R., WILLIAMS T.A. and EVANS S.A. Breeding and evaluation of new white clover varieties for persistency and higher yields under grazing 403
EVANS S.A. *see* EVANS D.R.

FISHER A. *see* WILMAN D.
FISHER G.E.J., BAKER L.J. and. TILEY G.E.D. Herbage production from swards containing a range of grass, forb and clover species and under extensive management 58
FISHER G.E.J. and DOWDESWELL A.M. The effects of regrowth and maintenance height on a grass sward with a high density of tillers 464
FISHER G.E.J., DOWDESWELL A.M. and PERROTT G. The effects of sward characteristics and supplement type on the herbage intake and milk production of summer-calving cows 121
FITZGERALD J.J. Grass silage as a basic feed for store lambs. 1. Effect of wilting, chop length and stage of maturity of grass silage on intake and performance of store lambs 363
FITZGERALD J.J. Grass silage as a basic feed for store lambs. 2. Effect of harvesting system and chop length of grass silage on silage intake and performance of store lambs 378
FITZGERALD J.J. Grass silage as a basic feed for store lambs. 3. Effect of barley supplementation of silages varying in chop length on silage intake and lamb performance 389
FOTHERGILL M. *see* COLLINS R.P.
FRAME J. and MERRILEES D.W. The effect of tractor wheel passes on herbage production from diploid and tetraploid ryegrass swards 13
FRAME J. *see* GOODING R.F.
FRANKLIN M.F. *see* WIGHTMAN P.S.
FREDRICKSON E.L. *see* ESTELL R.E.

GALE G.E. *see* KNIGHT A.C.
GANDOIN J.M. *see* VOLAIRE F.
GILLILAND T.J. Response of perennial ryegrass varieties to annual changes in sward management 442
GONZALEZ-MURUA C. *see* ESTAVILLO J.M.
GOODING R.F., FRAME J. and THOMAS C. Effects of sward type and rest periods from sheep grazing on white clover presence in perennial ryegrass/white clover associations 180
GUPTA M. L. *see* AKKHARATH I.

HAIGH P.M., CHAPPLE D.G. and POWELL T.L. Effect of silage additives on big-bale grass silage 318
HARD D. L. *see* PHIPPS R.H.
HARRINGTON D. *see* KEADY T.W.J.
HAVSTAD K.M. *see* ESTELL R.E.
HEPP C., MILNE J.A., ILLIUS A. W. and BEATTIE M.M. The effect of summer management of perennial ryegrass-dominant swards on plant and animal responses in the autumn when grazed by sheep. 2. Herbage intake and grazing behaviour 260
HEPP C., MILNE J.A., ILLIUS A.W. and ROBERTSON E. The effect of summer management of perennial ryegrass-dominant swards on plant and animal

© 1996 Blackwell Science Ltd.

Grass and Forage Science (1996) Volume 51, 470–472

Author Index

ADRIAENS F. *see* PHIPPS R.H.
AKKHARATH I., GUPTA M.L. and TULLBERG J.N. The influence of weather on the effectiveness of mechanical and chemical conditioning on drying rate of lucerne hay 96

BAKER L.J. *see* FISHER G.E.J.
BAKEWELL E.L. *see* DAVIES D.R.
BARDGETT R. *see* DAVIES A.
BEATTIE M.M. *see* HEPP C.
BESGA G. *see* ESTAVILLO J.M.
BINNIE R.C., JOHNSTON D.T. and CHESTNUTT D.M.B. The effect of a high-magnesium perennial ryegrass variety on the magnesium status of sheep 456
BRIGNALL P. *see* DAVIES A.
BUCKINGHAM H. *see* SMITH R.S.
BULLARD M.J. *see* SMITH R.S.

CHAPPLE D.G. *see* HAIGH P.M.
CHESTNUTT D.M.B. *see* BINNIE R.C.
CLAWSON S. *see* MOWAT D.J.
COLGROVE P.M. *see* DEL POZO M.
COLLINS R.P., FOTHERGILL M. and RHODES I. Interactions between seedlings of perennial ryegrass and white clover cultivars in establishing swards 163
COLLINS R.P., FOTHERGILL M., POTTER J.F. and RHODES I. The use of repeated-measurements analysis to describe the overwintering of white clover cultivars 116
CORKHILL P. *see* SMITH R.S.
CUSHNAHAN A. *see* MCKEE C.A.

DAVIES A., SHAO J., BRIGNALL P., BARDGETT R., PARRY M. and POLLOCK C.J. Specification of climatic sensitivity of forage maize to climate change 306
DAVIES D.R., MERRY R.J. and BAKEWELL E.L. The effect of timing of slurry application on the microflora of grass, and changes occurring during silage fermentation 42
DE KERCHOVE G. *see* PHIPPS R.H.
DEL POZO M., WRIGHT I.A., WHYTE T.K. and COLGROVE P.M. Effects of grazing by sheep or goats on sward composition in ryegrass/white clover pasture and on subsequent performance of weaned lambs 142
DOWDESWELL A.M. *see* FISHER G.E.J.

EDWARDS G.R., PARSONS A.J., NEWMAN J.A. and WRIGHT I.A. The spatial pattern of vegetation in cut and grazed grass/white clover pastures 227, 219
ELBERSON H.W. and WEST C.P. Growth and water relations of field-grown tall fescue as influenced by drought and endophyte 333
ESTAVILLO J.M., GONZALEZ-MURUA C., BESGA G. and RODRIGUEZ M. Effect of cow slurry N on herbage productivity, efficiency of N utilization and on white clover content in a natural sward in the Basque Country, Spain 1
ESTELL R.E., FREDRICKSON E.L. and HAVSTAD K.M. Chemical composition of *Flourensia cernua* at four growth stages 434

EVANS D.R., WILLIAMS T.A. and EVANS S.A. Breeding and evaluation of new white clover varieties for persistency and higher yields under grazing 403
EVANS S.A. *see* EVANS D.R.

FISHER A. *see* WILMAN D.
FISHER G.E.J., BAKER L.J. and. TILEY G.E.D. Herbage production from swards containing a range of grass, forb and clover species and under extensive management 58
FISHER G.E.J. and DOWDESWELL A.M. The effects of regrowth and maintenance height on a grass sward with a high density of tillers 464
FISHER G.E.J., DOWDESWELL A.M. and PERROTT G. The effects of sward characteristics and supplement type on the herbage intake and milk production of summer-calving cows 121
FITZGERALD J.J. Grass silage as a basic feed for store lambs. 1. Effect of wilting, chop length and stage of maturity of grass silage on intake and performance of store lambs 363
FITZGERALD J.J. Grass silage as a basic feed for store lambs. 2. Effect of harvesting system and chop length of grass silage on silage intake and performance of store lambs 378
FITZGERALD J.J. Grass silage as a basic feed for store lambs. 3. Effect of barley supplementation of silages varying in chop length on silage intake and lamb performance 389
FOTHERGILL M. *see* COLLINS R.P.
FRAME J. and MERRILEES D.W. The effect of tractor wheel passes on herbage production from diploid and tetraploid ryegrass swards 13
FRAME J. *see* GOODING R.F.
FRANKLIN M.F. *see* WIGHTMAN P.S.
FREDRICKSON E.L. *see* ESTELL R.E.

GALE G.E. *see* KNIGHT A.C.
GANDOIN J.M. *see* VOLAIRE F.
GILLILAND T.J. Response of perennial ryegrass varieties to annual changes in sward management 442
GONZALEZ-MURUA C. *see* ESTAVILLO J.M.
GOODING R.F., FRAME J. and THOMAS C. Effects of sward type and rest periods from sheep grazing on white clover presence in perennial ryegrass/white clover associations 180
GUPTA M. L. *see* AKKHARATH I.

HAIGH P.M., CHAPPLE D.G. and POWELL T.L. Effect of silage additives on big-bale grass silage 318
HARD D. L. *see* PHIPPS R.H.
HARRINGTON D. *see* KEADY T.W.J.
HAVSTAD K.M. *see* ESTELL R.E.
HEPP C., MILNE J.A., ILLIUS A. W. and BEATTIE M.M. The effect of summer management of perennial ryegrass-dominant swards on plant and animal responses in the autumn when grazed by sheep. 2. Herbage intake and grazing behaviour 260
HEPP C., MILNE J.A., ILLIUS A.W. and ROBERTSON E. The effect of summer management of perennial ryegrass-dominant swards on plant and animal

© 1996 Blackwell Science Ltd.

responses in the autumn when grazed by sheep. 1. Tissue turnover and sward structure 250
HETHERINGTON R.A. *see* WRIGHT I.A.
HIRATA M. A new technique to describe canopy characteristics of grass swards with spatial distribution, dry-matter digestibility and dry weight of small-size canopy components 209
HOPKINS A., MARTYN T.M., JOHNSON R.H., SHELDRICK R.D. and LAVENDER R.H. Forage production by two *Lotus* species as influenced by companion grass species
HUNTER E.A. *see* WRIGHT I.A.

ILLIUS A.W. *see* HEPP C.

JARVIS S.C., WILKINS R.J. and PAIN B.F. Opportunities for reducing the environmental impact of dairy farming managements: a systems approach 21
JOHNSON R.H. *see* HOPKINS A.
JOHNSON R.H. *see* YARROW N.H.
JOHNSTON D.T. *see* BINNIE R.C.

KEADY T.W.J. and MURPHY J. Effects of inoculant treatment on ryegrass silage fermentation, digestibility, rumen fermentation, intake and performance of lactating dairy cattle 232
KEADY T.W.J. and O'KIELY P. An evaluation of the effects of rate of nitrogen fertilization of grassland on silage fermentation, in-silo losses, effluent production and aerobic stability 350
KEADY T.W.J. and STEEN R.W.J. Effects of applying a bacterial inoculant to silage immediately before feeding on silage intake, digestibility, degradability and rumen volatile fatty acid concentrations in growing beef cattle 155
KEADY T.W.J., MURPHY J.J. and HARRINGTON D. The effects of ensiling on dry-matter intake and milk production by lactating dairy cattle given forage as the sole feed 131
KNIGHT A.C., PRICE J.S. and GALE G.E. Mechanical removal of surface moistures from grass 412

LAIDLAW A.S. *see* TEUBER N.
LAVENDER R.H. *see* HOPKINS A.
LAWS J.A. *see* MISSELBROOK T.H.

MARINISSEN M.J. *see* WARRINGA J.W.
MARTYN T.M. *see* HOPKINS A.
MAXWELL A.J.F. *see* WRIGHT I.A.
MAYNE C.S. *see* MCKEE C.A.
MCKEE C.A., CUSHNAHAN A., MAYNE C.S. and UNSWORTH E.F. The effect of partial replacement of grass silage with fresh grass on rumen fermentation characteristics and rumen outflow rates in cattle 32
MERCHANT M. The intake of grass and rush (*Juncus effusus* L.) by goats grazing rush-infested grass pasture 81
MEREDITH R.H. and WARBOYS I.B. The use of roll-conditioning and potassium carbonate (K_2CO_3) to increase the drying rate of lucerne (*Medicago sativa* L.) 8
MERRILEES D.W. *see* FRAME J.
MERRY R.J. *see* DAVIES D.R.
MILLARD P. *see* THORTON B.
MILLWARD D. *see* SMITH R.S.
MILNE J.A. *see* HEPP C.
MISSELBROOK T.H., LAWS J.A. and PAIN B. Surface application and shallow injection of cattle slurry on grassland: nitrogen losses, herbage yields and nitrogen recoveries 270
MOWAT D.J. and CLAWSON S. Oviposition and hatching of the clover weevil *Sitona lepidus Gyll.* (Coleoptera: Curculionidae)
MURPHY J. *see* KEADY T.W.J.

NEWMAN J.A. *see* EDWARDS G.R.

O'KIELY P. *see* KEADY T.W.J.

PAIN B. *see* MISSELBROOK T.H.
PAIN B.F. *see* JARVIS S.C.
PARES D. *see* ROYO C.
PARRY M. *see* DAVIES A.
PARSONS A.J. *see* EDWARDS G.R
PENNING P. D. *see* YARROW N.H.
PERROTT G. *see* FISHER G.E.J.
PHIPPS R.H., ADRIAENS F., HARD D.L., DE KERCHOVE G. and WELLER R.F. Use of bovine somatotropin for milk production: the effect of time of treatment initiation on milk production of Holstein/ Friesian dairy cows placed on a high-forage, low-concentrate feeding system 111
POLLOCK C.J. *see* DAVIES A.
POTTER J.F. *see* COLLINS R.P.
POWELL T.L. *see* HAIGH P.M.
PRICE J.S. *see* KNIGHT A.C.

RAMMER C. Quality of grass silage infected with spores of *Clostridium tyrobutyricum* 88
RHODES I. *see* COLLINS R.P.
ROBERTSON E. *see* HEPP C.
RODRIGUEZ M. *see* ESTAVILLO J.M.
ROMAGOSA I. *see* ROYO C.
ROYO C. and PARES D. Yield and quality of winter and spring triticales for forage and grain 449
ROYO C. and ROMAGOSA I. Effect of forage removal at the first detectable node stage on the growth of winter and spring triticale 170
RUSSEL A.J.F. *see* WRIGHT I.A.

SHAO J. *see* DAVIES A.
SHELDRICK R.D. *see* HOPKINS A.
SHIEL R.S. *see* SMITH R.S.
SIBBALD A.R. *see* WRIGHT I.A.
SMITH R.S., BUCKINGHAM H., BULLARD M.J., SHIEL R.S. and YOUNGER A. The conservation management of mesotrophic (meadow) grassland in northern England. 1. Effects of grazing, cutting date and fertilizer on the vegetation of a traditionally managed sward 278
SMITH R.S., CORKHILL P., SHIEL R.S. and MILLWARD D. The conservation management of mesotrophic (meadow) grassland in northern England. 2. Effects of grazing, cutting date, fertilizer and seed application on the vegetation of an agriculturally improved sward 292
STEEN R.W.J *see* KEADY T.W.J.

TEUBER N. and LAIDLAW A.S. Influence of irradiance on branch growth of white clover stolons in rejected areas within grazed swards 73
THOMAS C. *see* GOODING R.F.
THORTON B. and MILLARD P. Nitrogen uptake by grasses: changes induced by competing neighbour plants differing in frequency of defoliation 242
TILEY G.E.D. *see* FISHER G.E.J.
TULLBERG J.N. *see* AKKHARATH I.

UNSWORTH E.F. *see* MCKEE C.A.

VOLAIRE F. and GANDOIN J.M. The effect of age of the sward on the relationship between water-soluble carbohydrate accumulation and drought survival in two contrasted populations of cocksfoot (*Dactylis glomerata* L.) 190

WARBOYS I.B. *see* MEREDITH R.H.

WARRINGA J.W. and MARINISSEN M.J. The effect of light intensity after anthesis on dry matter distribution and seed yield of *Lolium perenne* L. 103
WELLER R.F. *see* PHIPPS R.H.
WEST C.P. *see* ELBERSON H.W.
WHYTE T.K. *see* DEL POZO M.
WHYTE T.K. *see* WRIGHT I.A.
WIGHTMAN P.S., FRANKLIN M.F. and YOUNIE D. The response of mini-swards of perennial ryegrass/white clover to simulated rainfall following slurry application 325
WILKINS R.J. *see* JARVIS S.C.
WILLIAMS T.A. *see* EVANS D.R.
WILMAN D. and FISHER A. Effects of interval between harvests and application of fertilizer N in spring on the growth of perennial ryegrass in a grass/white clover sward 52
WRIGHT I.A. *see* DEL POZO M.
WRIGHT I.A. *see* EDWARDS G.R.
WRIGHT I.A., MAXWELL T.J., RUSSEL A.J.F., HUNTER E.A., SIBBALD A.R., HETHERINGTON R.A. and WHYTE T.K. The effect of grazed sward height and stocking rate on animal performance and output from beef cow systems 199

YARROW N.H., PENNING P. D. and JOHNSON R.H. The effect of plane winter nutrition and sward height on the performance of steers grazing grass/white clover swards 424
YOUNGER A. *see* SMITH R.S.
YOUNIE D. *see* WIGHTMAN P.S.